

# Clinical and Imaging Factors Associated with Lower Peak Oxygen Consumption in Patients with Hypertrophic Cardiomyopathy: the Value of Ventricular Strain by Magnetic Resonance Imaging

*Factores clínicos e imagenológicos asociados a un menor consumo máximo de oxígeno en pacientes con miocardiopatía hipertrófica: el valor del strain ventricular por resonancia magnética*

FLORENCIA B. CANTORA<sup>1</sup>, SANTIAGO DECOTTO<sup>1</sup>, SANTIAGO L. DEL CASTILLO<sup>1</sup>, ROCÍO BLANCO<sup>1</sup>, MARIANO L. FALCONI<sup>1</sup>, RODOLFO PIZARRO<sup>1</sup>, DIEGO PEREZ DE ARENAZA<sup>1</sup>

## ABSTRACT

**Background:** Peak oxygen consumption (peak  $VO_2$ ) is a key marker for assessing functional capacity and prognosis in patients with hypertrophic cardiomyopathy (HCM). Although multiple factors can influence this parameter, its relative impact and predictive value are not fully established.

**Objective:** The aim of this study was to analyze the association between clinical, echocardiographic, and cardiac magnetic resonance imaging (CMR) variables with peak  $VO_2$  in patients diagnosed with HCM.

**Methods:** A retrospective, observational, single-center cohort study was conducted in patients diagnosed with sarcomeric HCM belonging to an institutional HCM registry between January 2017 and March 2025, who had echocardiography, CMR, and oxygen consumption tests performed within less than a year of each other. Clinical, imaging, and functional data were collected from electronic medical records. Cardiac magnetic resonance scans were done with a 1.5-T magnet (Avanto, Siemens Medical Solutions®, Erlangen, Germany). Late gadolinium enhancement (LGE) quantification and global longitudinal strain by CMR (GLS-CMR) were obtained using Circle Cardiovascular Imaging software (Tissue Tracking, cvi42). Univariate and multivariate linear regression models were used to evaluate associations with peak  $VO_2$ .

**Results:** Fifty-four patients with mean age of  $53 \pm 18$  years (59% men) were included in the study. Mean peak  $VO_2$  was  $23.5 \pm 9.6$  mL/kg/min. In the univariate analysis, male sex ( $p=0.001$ ), indexed right ventricular end-diastolic volume ( $p<0.001$ ), and GLS-CMR ( $p=0.030$ ) were significantly associated with higher peak  $VO_2$ . Age was inversely associated with peak  $VO_2$  ( $p<0.001$ ). No significant associations were found with left ventricular ejection fraction, intraventricular obstructive gradient, left ventricular mass index, or LGE. In the multivariate analysis, the variables that showed an independent association with lower peak  $VO_2$  were female sex ( $p=0.007$ ), older age at diagnosis ( $p<0.001$ ), and lower GLS-CMR value ( $p=0.033$ ).

**Conclusions:** In patients with HCM, female sex, older age, and lower left ventricular global longitudinal strain by CMR were independently associated with lower peak  $VO_2$ . These findings highlight the usefulness of myocardial strain as a complementary functional marker that could contribute to improve prognostic stratification in this population.

**Key words:** Hypertrophic cardiomyopathy – Strain - Cardiac magnetic resonance imaging - Oxygen consumption

## RESUMEN

**Introducción:** El consumo pico de oxígeno ( $VO_2$  pico) es un marcador clave para evaluar la capacidad funcional y el pronóstico en pacientes con miocardiopatía hipertrófica (MCH). Aunque múltiples factores pueden influir en este parámetro, su impacto relativo y valor predictivo no están completamente establecidos.

**Objetivo:** Analizar la asociación entre variables clínicas, ecocardiográficas y de resonancia magnética cardíaca (RMC) con el  $VO_2$  pico en pacientes diagnosticados con MCH.

**Material y métodos:** Estudio retrospectivo, observacional y unicéntrico que incluyó pacientes con diagnóstico de MCH sarcomérica pertenecientes a un registro institucional entre 2017 y 2025, que contaran con ecocardiograma, RMC y prueba de consumo de oxígeno realizados en un intervalo menor a un año de diferencia entre sí. Se recolectaron de la historia clínica electrónica datos clínicos, imagenológicos y funcionales. Las exploraciones de RMC se realizaron con un imán de 1.5-T (Avanto, Siemens Medical Solutions®,

REV ARGENT CARDIOL 2026;94:xxx-xxx. <https://doi.org/10.7775/rac.v94.i1.20971>

SEE RELATED ARTICLE: REV ARGENT CARDIOL 2026;94:xxx-xxx. <https://doi.org/10.7775/rac.v94.i1.20981>

Received: 12/09/2025 – Accepted: 01/21/2026

**Correspondence:** Florencia Cantora. Department of Cardiology, Hospital Italiano de Buenos Aires. Argentina. Perón 4190. Postal code: C1199. Email: florencia.cantora@hospitalitaliano.org.ar.



<https://creativecommons.org/licenses/by-nc-sa/4.0/>

©Revista Argentina de Cardiología

<sup>1</sup> Department of Cardiology, Hospital Italiano de Buenos Aires., Autonomous City of Buenos Aires, Argentina.

Erlangen, Alemania). La cuantificación de realce tardío de gadolinio (RTG) y el *strain* longitudinal global por RMC (SLG-RMC) se efectuaron con el software Circle Cardiovascular Imaging (Tissue Tracking, cvi42). Se utilizaron modelos de regresión lineal univariados y multivariados para evaluar las asociaciones con el  $VO_2$  pico.

**Resultados:** Se incluyeron 54 pacientes con una edad promedio de  $53 \pm 18$  años, de los cuales el 59% eran hombres. El  $VO_2$  pico promedio fue de  $23,5 \pm 9,6$  mL/kg/min. En el análisis univariado, el sexo masculino ( $p=0,001$ ), el volumen telediastólico del ventrículo derecho indexado ( $p<0,001$ ) y el SLG-RMC ( $p=0,030$ ) se asociaron significativamente con un mayor  $VO_2$  pico. La edad presentó una asociación inversa con el  $VO_2$  pico ( $p<0,001$ ). No se hallaron asociaciones significativas con la fracción de eyección del ventrículo izquierdo, el gradiente obstructivo intraventricular, el índice de masa ventricular izquierda ni con el realce tardío de gadolinio. En el análisis multivariado, las variables que mostraron asociación independiente con un  $VO_2$  pico más bajo fueron el sexo femenino ( $p=0,007$ ), una mayor edad al diagnóstico ( $p<0,001$ ) y un menor valor de SLG-RMC ( $p=0,033$ ).

**Conclusiones:** En pacientes con MCH, el sexo femenino, una mayor edad y un menor *strain* longitudinal global del ventrículo izquierdo por RMC se asociaron de forma independiente con un menor  $VO_2$  pico. Estos hallazgos resaltan la utilidad del *strain* miocárdico como marcador funcional complementario que podría contribuir a mejorar la estratificación pronóstica en esta población.

**Palabras clave:** Miocardiopatía hipertrófica - *Strain* - Resonancia magnética cardíaca - Consumo de oxígeno

## INTRODUCTION

Hypertrophic cardiomyopathy (HCM) is a relatively common genetic clinical syndrome characterized by left ventricular (LV) hypertrophy in the absence of other cardiac, systemic, or metabolic disease (1). Its origin mainly lies in gene mutations of the sarcomere, and it is inherited in an autosomal dominant manner in most cases. (2) It presents a broad clinical spectrum, ranging from asymptomatic patients to those with advanced heart failure or sudden death. (3)

However, prognostic stratification in HCM remains challenging, given the heterogeneity in disease progression and the limited ability of conventional parameters to predict functional decline (3-5). Peak oxygen consumption (peak  $VO_2$ ), obtained through cardiopulmonary exercise testing, is a robust marker of functional capacity and an independent predictor of morbidity and mortality in various heart diseases, including HCM (6). Thus, peak  $VO_2$  was proposed as the primary endpoint in the main studies on this condition. (7-9) However, the clinical and imaging variables that determine lower peak  $VO_2$  in this population are not clearly defined.

Cardiac magnetic resonance imaging (CMR) has become the gold standard for LV morphological and functional assessment, also allowing accurate quantification of fibrosis using late gadolinium enhancement (LGE) (1,2). Recently, the incorporation of global longitudinal strain analysis by cardiac magnetic resonance imaging (GLS-CMR) through feature tracking detects earlier stages of ventricular involvement, which are not evident with conventional techniques such as volumetric left ventricular ejection fraction (LVEF) measurement (10, 11). It should be noted that GLS was used in this study because it is the most reproducible parameter and the most widely used in clinical practice, with less variability than radial or circumferential strain (12, 13).

However, the prognostic value of this parameter in relation to functional capacity in HCM still requires further evidence. (14-17)

## OBJECTIVES

We evaluated the association between GLS-CMR and other clinical, echocardiographic, and CMR variables

with peak  $VO_2$  measured in the cardiopulmonary exercise test in a contemporary cohort of patients with confirmed diagnosis of HCM, with the aim of identifying independent predictors of functional limitation that would optimize prognostic stratification in this group of patients.

## METHODS

A retrospective, observational, single-center cohort study was conducted including patients belonging to an institutional HCM registry between January 2017 and March 2025. All patients gave informed consent prior to inclusion in the study. The data for the present analysis were obtained from a review of the institutional electronic medical records.

Hypertrophic cardiomyopathy was defined as the presence of increased thickness in any LV segment  $>15$  mm, or  $>13$  mm in first-degree relatives diagnosed with this disease, in the absence of any other justifiable cause. (1) Whenever possible, genetic studies were performed to confirm the sarcomeric etiology. Patients over 18 years of age with a confirmed diagnosis of sarcomeric HCM were included in the study, and phenocopies or other causes of ventricular hypertrophy were excluded.

For this study, only those patients who had echocardiogram, CMR, and cardiopulmonary exercise testing performed within an interval  $\leq 12$  months from each other were selected.

The following variables were collected: demographic (age, sex, height), clinical (NYHA functional class, treatment received), echocardiographic (LVEF, maximum intraventricular gradient, presence of obstructive HCM defined by a gradient  $\geq 30$  mmHg at rest and/or systolic anterior movement of the anterior mitral valve), and CMR-derived parameters (indexed ventricular volumes and mass, LVEF, and right ventricular ejection fraction, maximum wall thickness, presence and quantification of LGE, and LV GLS-CMR).

Cardiac magnetic resonance imaging was performed using a 1.5 T scanner (Avanto, Siemens Medical Solutions®, Erlangen, Germany) and LGE quantification (using a 5-threshold technique) and GLS-CMR were acquired using Circle Cardiovascular Imaging software (Tissue Tracking, cvi42, Figure 1). Peak  $VO_2$  was obtained by cardiopulmonary testing on a cycle ergometer using an incremental protocol, expressed in mL/kg/min.

## Statistical analysis

Continuous variables were expressed as mean  $\pm$  standard deviation (SD) or as median and interquartile range (IQR),

depending on their distribution as assessed by the Shapiro-Wilk test, and were compared using Student's t-test or the Mann-Whitney test, depending on data distribution. Categorical variables were presented as absolute frequencies and percentages and were compared using the Chi<sup>2</sup> test or Fisher's exact test, as appropriate.

Linear regression models were applied to explore the associations between clinical, echocardiographic, and CMR variables with peak VO<sub>2</sub>. In the univariate analysis, each predictor was evaluated individually to identify significant correlations with peak VO<sub>2</sub>. Subsequently, those variables with statistical significance ( $p < 0.05$ ) or clinical relevance were incorporated into a multivariate linear regression model to identify independent predictors of lower peak VO<sub>2</sub>. All statistical analyses were performed using STATA software version 13.1 (StataCorp LP, College Station, TX, USA), and  $p$  values  $< 0.05$  were considered as statistically significant.

#### Ethical considerations

The study protocol follows the ethical guidelines of the Declaration of Helsinki, (18) and the design of the institutional hypertrophic cardiomyopathy registry from which the data were obtained was approved by the institutional ethics committee.

#### RESULTS

A total of 54 patients diagnosed with sarcomeric HCM were included in the study. Mean age was  $53 \pm 18$  years, and 59.2% ( $n=32$ ) were men. The obstructive form of the disease was recorded in 26 pa-

tients (48.1%). Mean peak VO<sub>2</sub> in the population was  $23.5 \pm 9.6$  mL/kg/min.


#### Cardiac magnetic resonance parameters

Mean LVEF was  $72 \pm 10\%$  and mean maximum wall thickness  $17 \pm 4$  mm. Late gadolinium enhancement was present in 94.4% ( $n=51$ ) of patients, with a median mass of 15 grams and a 95% confidence interval (CI) of 6.84-33.59 (no discrimination was made according to LGE percentage). Mean LV GLS-CMR was  $-13 \pm 3.3\%$ . Figure 1 shows an example of GLS-CMR measurement in one cohort patient. The remaining clinical, echocardiographic, CMR, and cardiopulmonary exercise test variables are detailed in Table 1.

#### Univariate analysis

In the univariate linear regression model, significant associations were observed between peak VO<sub>2</sub> and age at diagnosis ( $\beta = -0.35$ ; 95% CI: -0.46 to -0.24;  $p < 0.001$ ), male sex ( $\beta = +8.26$ ; 95% CI +3.37 to +13.14;  $p = 0.001$ ), indexed right ventricular end-diastolic volume ( $\beta = +0.31$ ; 95% CI +0.15 to +0.46;  $p < 0.001$ ), and LV GLS-CMR ( $\beta = -0.85$ ; 95% CI -1.62 to -0.08;  $p = 0.034$ ). No significant associations were found with NYHA functional class, the presence of dynamic intraventricular obstruction, maximum intraventricular gradient, LVEF, LV volumes, LV mass, maximum wall thickness, or LGE. Table 2 details the variables

**Fig. 1.** Example of left ventricular global longitudinal strain analysis using Circle Cardiovascular Imaging (Tissue Tracking, cvi42).



**Table 1.** General characteristics of the population

Clinical variables	
Age at diagnosis, years, mean $\pm$ SD	53 $\pm$ 18
Male sex, n (%)	32 (59.3)
Height, centimeters, mean $\pm$ SD	161 $\pm$ 33
NYHA functional class* $\geq$ 2, n (%)	16 (29.6)
BMI*, kg/m <sup>2</sup> , mean $\pm$ SD	28.4 $\pm$ 5.1
Hypertension, n (%)	21 (38.9)
Diabetes mellitus, n (%)	6 (11.1)
Atrial fibrillation, n (%)	7 (13.0)
Dyspnea, n (%)	31 (57.4)
Angina, n (%)	12 (22.2)
Atrial fibrillation, n (%)	7 (13.0)
Beta-blockers, n (%)	38 (70.4)
Calcium channel blockers, n (%)	7 (13.0)
Myosin inhibitors, n (%)	3 (5.6)
ICD*, n (%)	6 (11.1)
Septal myectomy, n (%)	3 (5.6)
Septal alcoholization, n (%)	2 (3.7)
Echocardiographic variables	
Wall thickness, mm, mean $\pm$ SD	17.4 $\pm$ 3.9
Left atrial anteroposterior diameter, mm, median (IQR)	42 [38-46]
Intraventricular gradient, mmHg, median (IQR)	30 [6-58]
SAM, n (%)	25 (46.3)
LVOTO, n (%)	32 (59.3)
Cardiac resonance variables	
LVEF, %, mean $\pm$ SD	72 $\pm$ 10
Indexed LV end-diastolic volume, mL/m <sup>2</sup> , mean $\pm$ SD	72 $\pm$ 18
Indexed LV end-systolic volume, mL/m <sup>2</sup> , median (IQR)	69 [60-78]
Maximum thickness, mm, mean $\pm$ SD	17 $\pm$ 4
Presence of LGE, n (%)	51 (94.4)
LGE mass, grams, median (IQR)	15 [6.84-33.59]
Left ventricular GLS, %, mean $\pm$ SD	-13 $\pm$ 3.3
Right ventricular ejection fraction, %, mean $\pm$ SD	66 $\pm$ 7
RV end-diastolic volume, mL, mean $\pm$ SD	71 $\pm$ 15
RV end-systolic volume, mL, mean $\pm$ SD	27 $\pm$ 24
LV mass, grams	86 $\pm$ 24
Oxygen consumption variables	
Peak VO <sub>2</sub> (mL/kg/min), mean $\pm$ SD	23.5 $\pm$ 9.6
VO <sub>2</sub> (% of predicted), mean $\pm$ SD	83.3 $\pm$ 19.5
Heart rate at peak VO <sub>2</sub> (bpm), mean $\pm$ SD	133.7 $\pm$ 30.7
OUES, [(mL/min)/log10], mean $\pm$ SD	2156.8 $\pm$ 954.7

BMI: body mass index; ICD: implantable cardioverter defibrillator; IQR: interquartile range; LGE: late gadolinium enhancement; LV: left ventricular; LVEF: left ventricular ejection fraction; LVOTO: left ventricular outflow tract obstruction; NYHA: New York Heart Association; OUES: oxygen uptake efficiency slope; RV: right ventricular; SAM: systolic anterior mitral valve movement; SD: standard deviation.

included in the univariate analysis and their relationship with peak  $\text{VO}_2$ .

### Multivariate analysis

In the multivariate linear regression model (Table 3), the independent predictors of lower peak  $\text{VO}_2$  were: female sex ( $\beta = -6.1$ ; 95% CI -10.56 to -1.75;  $p = 0.007$ ), older age at diagnosis ( $\beta = -0.26$ ; 95% CI: -0.37 to -0.15;  $p < 0.001$ ), and lower LV GLS-CMR ( $\beta = -0.59$ , 95% CI: -1.14 to -0.04;  $p = 0.033$ ).

### DISCUSSION

In this contemporary cohort of patients with HCM, we observed that a lower LV-GLS-CMR value was independently associated with lower functional capacity, as measured by peak  $\text{VO}_2$  in the cardiopulmonary exercise test. This finding could provide a relevant pathophysiological perspective, based on the hypoth-

esis that functional limitation in HCM would depend not only on the degree of hypertrophy or the presence of dynamic obstruction, but also on both clinical and subclinical myocardial contractile failure, detectable by GLS-CMR, even in patients with preserved LVEF. Global longitudinal strain, by reflecting the longitudinal deformation capacity of the myocardium, is a sensitive marker of contractile efficiency and allows the detection of early mechanical dysfunction before global LVEF alterations or advanced symptoms become apparent (19, 20). It is noteworthy that all patients in our cohort had preserved LVEF.

Abnormal GLS probably represents the functional expression of the distinctive molecular mechanisms of HCM, characterized by a progressive loss of contractile efficiency, which translates into reduced myocardial shortening and, consequently, a limited ability to increase cardiac output during exercise. In other words,

**Table 2.** Univariate linear regression model

Variable	Univariate linear regression $\beta$ coefficient (95% CI)	p value
Age at diagnosis	-0.35 (-0.46 to -0.24)	<0.001
Male	+8.26 (+3.37 to +13.14)	0.001
NYHA dyspnea	-2.59 (-5.35 to +0.17)	0.065
Maximum intraventricular gradient (mmHg)	-0.06 (-0.15 to +0.02)	0.110
Obstructive HCM	-1.14 (-6.44 to +4.17)	0.669
LVEF %	+0.11 (-0.16 to +0.38)	0.425
Indexed left ventricular end-diastolic volume, (mL/m <sup>2</sup> )	+0.08 (-0.06 to +0.23)	0.265
Indexed LV end-systolic volume, (mL/m <sup>2</sup> )	-0.05 (-0.15 to +0.05)	0.320
Left ventricular mass (g/m <sup>2</sup> )	-0.0007 (-0.11 to +0.11)	0.921
Maximum wall thickness (mm)	+0.09 (-0.60 to +0.79)	0.781
Quantitative LGE (%)	-0.004 (-0.10 to +0.09)	0.934
LGE (present)	+2.35 (-18 to 23.2)	0.821
RVEF (%)	+0.25 (-0.10 to +0.60)	0.162
Indexed RV end-diastolic volume (mL/m <sup>2</sup> )	+0.31 (+0.15 to +0.46)	<0.001
Indexed RV end-systolic volume (mL/m <sup>2</sup> )	-0.018 (-0.13 to +0.09)	0.743
CMR strain (%)	-0.85 (-1.6 to -0.08)	0.034

CMR: cardiac magnetic resonance; HCM: hypertrophic cardiomyopathy; LGE: late gadolinium enhancement; LV: left ventricular; LVEF: left ventricular ejection fraction; NYHA: New York Heart Association; RV: right ventricular; RVEF: right ventricular ejection fraction.

**Table 3.** Multivariate linear regression model

Variable	Multivariate linear regression $\beta$ coefficient (95% CI)	p value
Age at diagnosis (years)	-0.26 (-0.37 to -0.15)	<0.001
Female sex	-6.10 (-10.56 to -1.75)	0.007
LV global longitudinal strain	-0.59 (-1.14 to -0.04)	0.033

LV: left ventricular

reduced GLS implies a lower contractile reserve and, therefore, a limitation in sustaining cardiac output, which translates into decreased peak  $\text{VO}_2$ . (14,19) This phenomenon could explain the association between greater GLS-CMR alteration and lower peak  $\text{VO}_2$  observed in our study, in accordance with the findings of the STRAIN-HCM study, where GLS behaved as an independent predictor of adverse events. (14, 21)

On the other hand, we found no correlation between peak  $\text{VO}_2$  and the amount of myocardial fibrosis measured by LGE, a finding that reinforces the idea that GLS identifies an earlier functional stage of myocardial damage. (22, 23) It should be noted that in our cohort, almost all patients had some degree of LGE, suggesting that the differential prognostic value of GLS does not depend exclusively on the presence or absence of fibrosis, but on the degree of underlying mechanical dysfunction.

The lower functional capacity observed in women in this study is consistent with reports from other series, where female sex is associated with more symptomatic phenotypes, smaller ventricular cavities, and greater predisposition to dynamic obstruction. (24) These anatomical features, combined with a frequently later diagnosis, could partly explain the functional gap observed. Similarly, older age was a negative determinant of peak  $\text{VO}_2$ , probably reflecting the cumulative impact of diffuse fibrosis, myocardial stiffness, and associated comorbidities. (25, 26)

From a clinical perspective, incorporating GLS-CMR into the evaluation of HCM may provide additional prognostic value and allow for a more accurate characterization of the functional phenotype. In patients with preserved LVEF, a significant reduction in GLS could anticipate clinical deterioration and guide early interventions. This integrated approach is in line with the current trend in the AHA/ACC international guidelines, (1) which promote personalized pathophysiological stratification, prevailing over a purely morphological approach.

In the future, it will likely be necessary to develop prognostic models that integrate clinical, imaging, and functional variables to refine risk prediction in this population.

Although this study has limitations inherent to its retrospective design and the single-center nature of the sample, the availability of high-quality complementary studies—echocardiogram, CMR, and cardiopulmonary testing—strengthens the robustness of the results. Furthermore, our population represents a clinical profile characteristic of HCM, with the presence of obstructive forms, which reinforces the representativeness of the sample and the applicability of the findings. Taken together, our data consolidate the notion that HCM is essentially a disease of myocardial mechanics rather than hypertrophy per se, and position GLS-CMR as a potentially important marker for understanding, quantifying, and monitoring contractile efficiency in clinical practice.

## CONCLUSIONS

In this contemporary cohort of patients with HCM, female sex, older age, and reduced GLS-CMR were independently associated with lower peak  $\text{VO}_2$ .

These results highlight the usefulness of GLS-CMR as a complementary tool for prognostic stratification and functional assessment, beyond conventional parameters such as LVEF or LGE. Its systematic implementation could optimize the early identification of patients at higher risk of functional deterioration.

## Limitations

The limitations of this study include its retrospective nature and the lack of stratification of peak  $\text{VO}_2$  according to patient sex. It is also important to note that no comparison was made between strain measured by cardiac magnetic resonance imaging and strain assessed by echocardiography.

## Conflicts of interest

None declared.

(See authors' conflict of interests forms on the web).

## Acknowledgments

To the entire cardiology department of Hospital Italiano de Buenos Aires.

## Sources of funding

None.

## REFERENCES

1. Writing Committee Members; Ommen SR, Ho CY, Asif IM, Balaji S, Burke MA, Day SM, Dearani JA, et al. 2024 AHA/ACC/AMSSM/HRS/PACES/SCMR Guideline for the Management of Hypertrophic Cardiomyopathy: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol* 2024;83:2324-405. <https://doi.org/10.1016/j.jacc.2024.02.014>.
2. Ho CY, Day SM, Ashley EA, Michels M, Pereira AC, Jacoby D, et al. Genotype and lifetime burden of disease in hypertrophic cardiomyopathy: insights from the Sarcomeric Human Cardiomyopathy Registry (SHaRe). *Circulation* 2018;138:1387-98. <https://doi.org/10.1161/CIRCULATIONAHA.117.033200>
3. Marian AJ, Braunwald E. Hypertrophic cardiomyopathy: genetics, pathogenesis, clinical manifestations, diagnosis, and therapy. *Circ Res* 2017;121:749-70. <https://doi.org/10.1161/CIRCRESAHA.117.311059>
4. Fumagalli C, De Filippo V, Zocchi C, Tassetti L, Marra MP, Brunetti G, et al. Strength of clinical indication and therapeutic impact of the implantable cardioverter defibrillator in patients with hypertrophic cardiomyopathy. *Int J Cardiol* 2022;353:62-7. <https://doi.org/10.1161/CIRCRESAHA.117.311059>
5. Maron BJ, Spirito P, Ackerman MJ, Casey SA, Semsarian C, Estes NA 3rd, et al. Prevention of sudden cardiac death with implantable cardioverter-defibrillators in children and adolescents with hypertrophic cardiomyopathy. *J Am Coll Cardiol* 2013;61:1527-35. <https://doi.org/10.1016/j.jacc.2013.01.037>
6. Rowin EJ, Maron BJ, Olivetto I, Maron MS. Role of exercise testing in hypertrophic cardiomyopathy. *JACC Cardiovasc Imaging*. 2017;10:1374-86. <https://doi.org/10.1016/j.jcmg.2017.07.016>
7. Rossano BJ, Canter C, Wolf C, Favatella N, Lockman J, Puli S, et al. Mavacamten in symptomatic adolescent patients with obstructive hypertrophic cardiomyopathy: design of the phase 3 SCOUT-HCM trial. *Am Heart J* 2026;292:107283. <https://doi.org/10.1016/j.ahj.2025.107283>

8. Desai MY, Gaballa A, Okushi Y, Abusafia M, Jadam S, Rutkowski K, et al. Real-World Observations in Patients With Obstructive Hypertrophic Cardiomyopathy Treated With Mavacamten: Evidence of Favorable Disease Modification. *J Am Heart Assoc* 2025;14:e044537. <https://doi.org/10.1161/JAHA.125.044537>
9. Reddy Kalluru PK, Siddenti SM, Valisekka SS, Suddapalli SK, Juturu UT, Chagam Reddy S, et al. A Systematic Review of Clinical Trials on Mavacamten in Hypertrophic Cardiomyopathy. *Heart Int* 2025;19:31-40. <https://doi.org/10.17925/HI.2025.19.1.7>
10. Pezeshki PS, Ghorashi SM, Houshmand G, Ganjparvar M, Pouraliakbar H, Rezaei-Kalantari K, et al. Feature tracking cardiac magnetic resonance imaging to assess cardiac manifestations of systemic diseases. *Heart Fail Rev* 2023;28:1189-99. <https://doi.org/10.1007/s10741-023-10321-6>
11. Amzulescu MS, De Craene M, Langet H, Pasquet A, Vancraeynest D, Pouleur AC, et al. Myocardial strain imaging: review of general principles, validation, and sources of discrepancies. *Eur Heart J Cardiovasc Imaging* 2019;20:605–19. <https://doi.org/10.1093/ehjci/jez041>
12. Karlsen S, Dahlslett T, Grenne B, Sjøli B, Smiseth O, Edvardsen T, et al. Global longitudinal strain is a more reproducible measure of left ventricular function than ejection fraction regardless of echocardiographic training. *Cardiovasc Ultrasound* 2019;17:18. <https://doi.org/10.1186/s12947-019-0168-9>
13. Augustine D, Lewandowski AJ, Lazdam M, Rai A, Francis J, Myerson S, et al. Global and regional left ventricular myocardial deformation measures by magnetic resonance feature tracking in normal subjects: reproducibility and reference values. *J Cardiovasc Magn Reson* 2013;15:8. <https://doi.org/10.1186/1532-429X-15-8>
14. Negri F, Muser D, Driussi M, Sanna GD, Masè M, Cittar M, et al. Prognostic role of global longitudinal strain by feature tracking in patients with hypertrophic cardiomyopathy: The STRAIN-HCM study. *Int J Cardiol* 2021;345:61-7. <https://doi.org/10.1016/j.ijcard.2021.10.148>
15. Albulushi A, Abri QA, Tawfek A, Bagheri AR, Al-Hinai S. Review Article--Exercise and Hypertrophic Cardiomyopathy: Risks, Benefits, and Safety - A Systematic Review and Meta-analysis. *J Saudi Heart Assoc* 2025;37:9. <https://doi.org/10.37616/2212-5043.1421>
16. Scatteia A, Baritussio A, Bucciarelli-Ducci C. Strain imaging using cardiac magnetic resonance. *Heart Fail Rev* 2017;22:465-76. <https://doi.org/10.1007/s10741-017-9621-8>
17. Kawel-Boehm N, Hetzel SJ, Ambale-Venkatesh B, Captur G, Francois CJ, Jerosch-Herold M, et al. Reference ranges (“normal values”) for cardiovascular magnetic resonance in adults and children: 2020 update. *J Cardiovasc Magn Reson* 2020;22:87. <https://doi.org/10.1186/s12968-020-00683-3>
18. World Medical Association. World Medical Association Declaration of Helsinki: Ethical Principles for Medical Research Involving Human Subjects. *JAMA*. 2013;310 :2191–2194. doi:10.1001/jama.2013.281053
19. Romano S, Judd RM, Kim RJ, Kim HW, Klem I, Heitner JF, et al. Feature-Tracking Global Longitudinal Strain Predicts Death in a Multicenter Population of Patients With Ischemic and Nonischemic Dilated Cardiomyopathy Incremental to Ejection Fraction and Late Gadolinium Enhancement. *JACC Cardiovasc Imaging* 2018;11:1419-29. <https://doi.org/10.1016/j.jcmg.2017.10.024>
20. Canciello G, Lombardi R, Borrelli F, Ordine L, Chen SN, Santoro C, et al. Echocardiographic Strain Abnormalities Precede Left Ventricular Hypertrophy Development in Hypertrophic Cardiomyopathy Mutation Carriers. *Int J Mol Sci* 2024;25:8128. <https://doi.org/10.3390/ijms25158128>
21. Maron, M, Wells, S. Myocardial Strain in Hypertrophic Cardiomyopathy: A Force Worth Pursuing?. *J Am Coll Cardiol Img* 2019;12:1943-45. <https://doi.org/10.1016/j.jcmg.2018.09.026>
22. Saijo Y, Van Iterson E, Vega Brizneda M, Desai MY, Lever HM, Smedira NG, et al. Impact of left atrial strain mechanics on exercise intolerance and need for septal reduction therapy in hypertrophic cardiomyopathy. *Eur Heart J Cardiovasc Imaging* 2022;23:238-45. <https://doi.org/10.1093/ehjci/jeab001>
23. Tower-Rader A, Mohananey D, To A, Lever HM, Popovic ZB, Desai MY, et al. Prognostic Value of Global Longitudinal Strain in Hypertrophic Cardiomyopathy: A Systematic Review of Existing Literature. *J Am Coll Cardiol Img* 2019;12:1930-42. <https://doi.org/10.1016/j.jcmg.2018.07.016>
24. Trongtorsak A, Polpichai N, Thangjui S, Kewcharoen J, Yodsuan R, Devkota A, Friedman HJ, Estrada AQ. Gender-Related Differences in Hypertrophic Cardiomyopathy: A Systematic Review and Meta-Analysis. *Pulse (Basel)* 2021;9:38-46. <https://doi.org/10.1159/000517618>
25. Mehta LS, Beckie TM, DeVon HA, Grines CL, Krumholz HM, Johnson MN, et al. Acute myocardial infarction in women: a scientific statement from the American Heart Association. *Circulation* 2016;133:916–47. <https://doi.org/10.1161/CIR.0000000000000351>
26. Reant P, Mirabel M, Lloyd G, Peyrou J, Lopez Ayala JM, Dickie S, et al. Global longitudinal strain is associated with heart failure outcomes in hypertrophic cardiomyopathy. *Heart* 2016;102:741-7. <https://doi.org/10.1136/heartjnl-2015-308576>