

# Characteristics and Evolution of Cardiogenic Shock According to Gender in Latin America. LATIN Shock Registry Data

## *Características y evolución del shock cardiogénico de acuerdo con el sexo en Latinoamérica. Datos del registro LATIN Shock*

YANINA CASTILLO COSTA<sup>1, MTSAC</sup>, FLAVIO DELFINO<sup>2, MTSAC</sup>, JOSÉ MACÍAS<sup>2</sup>, MIGUEL QUINTANA<sup>3</sup>, MARIANO ADAMOWSKI<sup>4</sup>, FABIOLA RODRÍGUEZ CABALLERO<sup>1, 5</sup>, MA DE LOS ÁNGELES NAVARTA NAVARRO<sup>1, 6</sup>, GABRIELA VELAZCO<sup>1, 7</sup>, VÍCTOR MAURO<sup>1, MTSAC</sup>, ON BEHALF OF LATIN SHOCK GROUP

### ABSTRACT

**Background:** Cardiogenic shock (CS) as a complication of myocardial infarction is a condition with high mortality that, due to biological or equity differences, may have different characteristics according to gender.

**Objectives:** The aim of this study was to evaluate whether there are differences in the presentation, clinical evolution and treatment of patients with CS, according to gender.

**Methods:** We analyzed the patients of the LATIN Shock registry and evaluated whether there were differences according to gender.

**Results:** A total of 278 patients (30% women) were included. Seventy-five percent of patients presented ST-segment elevation acute coronary syndrome. Women were older and men were heavier smokers, and there were no differences in the prevalence of other cardiovascular risk factors, history, or comorbidities between genders. Revascularization was similar in men and women (86%) and there were no differences in the indication for pulmonary catheterization (11% vs. 20%,  $p=0.082$ ). Women received less mechanical support with intra-aortic balloon counterpulsation (14% vs. 26%,  $p=0.032$ ), while use of other mechanical support procedures was scarce, only in 1% of women and in 3% of men ( $p=0.678$ ). Mortality in women was 48% vs. 54% in men, not reaching statistical significance ( $p=0.470$ ).

**Conclusions:** Women with CS due to infarction are older and receive similar treatment except for mechanical supports, which are more frequently used in men. In our study there were no significant differences in mortality according to gender, which was high in both groups.

**Key words:** Cardiogenic Shock - Myocardial Infarction - Gender - Registry.

### RESUMEN

**Introducción:** El shock cardiogénico (SC) como complicación del infarto agudo de miocardio (IAM) es una patología con alta mortalidad que por diferencias biológicas o de equidad podría tener distintas características de acuerdo con el sexo.

**Objetivos:** Evaluar si hay diferencias en la presentación, evolución clínica y tratamiento de acuerdo con el sexo en los pacientes con SC.

**Material y métodos:** Se analizaron los pacientes del registro LATIN Shock y se evaluó si existían diferencias de acuerdo con el sexo.

**Resultados:** Se incluyeron 278 pacientes (30 % mujeres), el 75 % con síndrome coronario agudo con elevación del segmento ST (SCACEST). Las mujeres fueron más añosas y los hombres más frecuentemente tabaquistas. No hubo diferencias en la prevalencia de otros factores de riesgo cardiovascular, antecedentes ni comorbilidades entre ambos sexos. Hombres y mujeres recibieron revascularización en forma similar (86 %). No hubo diferencias en la indicación de cateterismo pulmonar (11 % vs 20 %,  $p = 0,082$ ). Las mujeres recibieron menos soporte mecánico con balón de contrapulsación (14 % vs 26 %,  $p=0,032$ ).

El empleo de otros soportes mecánicos fue escaso (1 % de las mujeres y 3 % de los hombres,  $p= 0,678$ ). La mortalidad de las mujeres fue del 49 % y la de los hombres 54 %, sin diferencia significativa ( $p= 0,470$ ).

REV ARGENT CARDIOL 2024;92:403-409. <http://dx.doi.org/10.7775/rac.v92.i6.20842>

SEE RELATED ARTICLE: REV ARGENT CARDIOL 2024;92:349-400. <http://dx.doi.org/10.7775/rac.v92.i6.20846>

Received: 10/12/2024 – Accepted: 12/06/2024

Correspondence: Yanina Castillo Costa. Azcuenaga 980. E-mail: [yanu\\_c@hotmail.com](mailto:yanu_c@hotmail.com)



<https://creativecommons.org/licenses/by-nc-sa/4.0/>

©Revista Argentina de Cardiología

<sup>1</sup> Clínica Bazterrica, Autonomous City of Buenos Aires, Argentina

<sup>2</sup> Clínica Santa Isabel, Autonomous City of Buenos Aires, Argentina

<sup>3</sup> Instituto Cardiovascular Migone, Asunción, Paraguay

<sup>4</sup> Hospital El cruce, Florencio Varela, Pcia. de Buenos Aires, Argentina

<sup>5</sup> Instituto Nacional de Cardiología Profesor Dr. Juan A Cattoni, Asunción, Paraguay

<sup>6</sup> Hospital descentralizado Dr. Guillermo Rawson, San Juan, Argentina

<sup>7</sup> Hospital Santojanni, Autonomous City of Buenos Aires, Argentina

**Conclusiones:** Las mujeres con SC por IAM son más añosas que los hombres y reciben similar tratamiento, excepto soportes mecánicos, más utilizados en hombres. No encontramos diferencias significativas de acuerdo con el sexo en la mortalidad, que fue elevada en ambos grupos.

**Palabras clave:** Shock cardiogénico - Infarto de miocardio - Sexo - Registro

## INTRODUCTION

Cardiovascular disease is the leading cause of death in both men and women. (1) It has been shown that in myocardial infarction, female gender, usually underrepresented in published works, (2,3) is associated with longer delays in consultation, lower rates of primary angioplasty, higher probability of developing heart failure and cardiogenic shock (CS), (4-6) and higher mortality. (7-10). In CS, a condition with mortality ranging between 40-60%, (11) differences in presentation, treatment received and evolution between men and women are not clearly defined. This "indefiniteness" is particularly relevant today, since there is evidence that biological gender (considering the different hormonal influence) implies some differences in physiological responses and in the action of certain drugs (12) that influence clinical disorders [as in sepsis (13, 14)], and should therefore be taken into special account. (1) Similarly, equity in access to health care is a current objective of many scientific societies. (15) To determine whether there is equity or differences in relation to gender, the contribution of regional multicenter registries, in our case the LATIN Shock registry, (16) is essential, since it has also been shown that there are differences in access to health care depending on whether high or low-income countries are evaluated. (17,18)

To date, there is scarce literature worldwide on the implications of gender differences in the characteristics, evolution and treatment of CS, and none in Latin America.

## OBJECTIVES

1) To analyze whether there are differences in the clinical presentation, treatment received, and clinical evolution according to gender in patients with CS in the context of acute coronary syndromes (ACS), and 2) To establish whether gender is an independent predictor of in-hospital mortality.

## METHODS

LATIN Shock is a multicenter, observational, prospective and consecutive registry of ACS complicated with CS. The Research Area of the Argentine Society of Cardiology provided computer and statistical support for the study.

Cardiogenic shock was defined as the presence of systolic blood pressure (SBP) <90 mmHg for at least 30 minutes or requirement of vasopressors and/or inotropic drugs to maintain SBP ≥90mmHg, associated with signs of hypoperfusion and signs of pulmonary congestion.

The inclusion criteria were: patients over 18 years of age, hospitalized in a coronary care unit or multipurpose critical

care unit for ST-segment elevation ACS (STE-ACS) or non-ST-segment elevation ACS (NSTEMI-ACS), who presented CS since admission or developed it during hospitalization.

Data were collected by the responsible investigators of the different centers in an electronic file designed ad hoc with the RedCAP platform. In-hospital events were analyzed.

Further details on the characteristics of the participating centers can be found in LATIN Shock. (16)

## Statistical analysis

The information obtained through RedCAP was exported in Excel and the database was analyzed using Epi-info 7. Continuous variables were expressed as mean and standard deviation for those with normal distribution and as median with interquartile range 25% -75% (IQR) for those non normal distribution. The statistical analysis of continuous variables was performed using Student's t test or the Wilcoxon rank sum test, as appropriate. Discrete variables were expressed as percentages and comparisons were performed using the chi-square test with Yates correction or Fisher's exact test, as appropriate.

Contingency tables were built to analyze variable association or independence. Linear and/or multiple logistic regression analyses were carried out to determine the existence of associations and/or independent predictions between the different variables involved and mortality. All statistical comparisons were two-tailed and values of  $p < 0.05$  were considered statistically significant.

## Ethical considerations


The study was conducted in accordance with current national and international regulations: Declaration of Helsinki of the World Medical Association, the ICH E6 Good Clinical Practice Guidelines, Resolution 1480/11 of the National Ministry of Health and GCBA Law 3301/09. The protocol was approved by the Bioethics Committee of the Argentine Society of Cardiology (SAC). Informed consent was waived since no sensitive data were recorded and only in-hospital follow-up was performed.

## RESULTS

Forty-one Latin American centers from seven countries (Argentina, Bolivia, Chile, Ecuador, Honduras, Paraguay and Perú) participated in the registry, recruiting 278 patients from October 2021 to September 2023. Seventy percent of patients were men ( $n=195$ ) and overall median age was 66 years (IQR 59-75). Women were older [median age 71 years (IQR 61-78) vs. 64 years (IQR 58-73) in men,  $p < 0.001$ ]. Most CS (75%) were due to STE-ACS. Details of the causes of CS are shown in Figure 1.

CS was present from admission in 60% of cases (66% in women and 57% in men;  $p=0.181$ ). In those admitted with another Killip, 81% developed CS on

Fig. 1. Causes of CS



CS: Cardiogenic shock; Mec Complic: mechanical complications; NSTEMI-ACS: non-ST-segment elevation acute coronary syndrome; RV AMI: right ventricular acute myocardial infarction; STE-ACS: ST-segment elevation acute coronary syndrome

Table 1. Baseline characteristics of the population.

	Total (n=278)	Female (n=83)	Male (n=195)	p
Age (years)	66 (59-75)	71 (61-78)	64 (58-73)	<0.001
Hypertension	206 (74)	66 (80)	140 (72)	0.179
Dyslipidemia	95 (34)	28 (34)	67 (34)	0.920
Smoking	72 (26)	13 (16)	59 (30)	0.011
Type 2 diabetes	117 (42)	35 (42)	82 (42)	0.494
Obesity	79 (29)	26 (31)	53 (27)	0.483
COPD	19 (7)	3 (4)	16 (8)	0.165
Chronic anemia	14 (5)	5 (6)	9 (5)	0.623
Chronic kidney disease	17 (6)	4 (5)	13 (7)	0.556
Previous AMI	44 (16)	8 (10)	36 (19)	0.065
Previous HF	20 (7)	6 (7)	14 (7)	0.483
Previous stroke	15 (5)	4 (5)	8 (4)	0.781

AMI: Acute myocardial infarction; COPD: Chronic obstructive pulmonary disease; HF: Heart failure. Quantitative variables are presented as median and interquartile range, qualitative variables as frequency and percentage.

the first day of hospitalization, 13% on the second day and 6% on the third day, with no differences according to gender.

Baseline characteristics according to gender are shown in Table 1

The use of vasoactive agents was 97.8%, with no differences by gender according to the type of drug used, except for vasopressin, which was used in 7.2% of women and 16% of men (p=0.041). Mechanical ventilation was used in 52.5% of cases in both genders, Swan Ganz catheter in 17% (11% in women and 20% in men, p=0.073), intra-aortic balloon counterpulsation in 22.2% (14% in women and 26% in men, p=0.032) and extracorporeal membrane oxygenation (ECMO) in 1% of women and 3% of men, p=0.676). No other supports were used.

The culprit vessel was revascularized in 81% of the overall population, with no differences according to sex (80% of women vs. 82% of men, p= 0.768).

The prevalence of multivessel disease was 71% (63% in women and 73% in men, p=0.113) and there was no difference in the prevalence of revascularization of other vessels, (40% in women and 31% in men, p=0.408).

In-hospital evolution by gender is detailed in Table 2.

The median (IQR) hospital stay was 6 days (1-16). Overall in-hospital mortality was 52.7%, with no differences between ACS with or without ST-segment elevation or according to gender: women 49% vs. men 54%, p=0.470).

The variables shown in Table 3 were considered for the analysis of univariate predictors of mortality

In the multivariate analysis model that included age, gender, diabetes, reperfusion and use of intra-aortic balloon counterpulsation, only age maintained its prognostic value (odds ratio 1.025, 95% CI 1.002-1.048, p=0.011).

In patients with STE-ACS without mechanical

complications on admission (n=208) there were no differences in baseline characteristics; median (IQR) time to overall consultation was 300 min (120-780); 360 min (120-540) in women and 274 min (120-870) in men, p=0.196. AMI was anterior in 68% of cases (women: 55.9% vs. men: 72.5%, p=0.012) and percutaneous coronary intervention (PCI) was performed in 88.9% of women and 82% of men (p=0.149) and the door-to-balloon time was 120 min (60-240), without differences according to gender. There were 19.1% of failed PCI in women and 21.2% in men (p=0.398). Mortality in women was 54.2% and 55.4% in men (p=0.437).

**DISCUSSION**

LATIN shock, the first Latin American study of CS in the context of ACS, allows us to learn about the reality of CS management in low-moderate income countries and at the same time, with a culture different from the European/American one, which could entail dis-

parities according to gender.

It is well known that the proportion of women with CS due to infarction is higher than that of men, as shown by the Argentine ARGEN IAM ST registry (6) or the French FAST AMI study, in which it is almost twice that of men. (19) This is probably due, among other things, to the fact that women have a longer life expectancy and, in general, are older at the time of infarction with shock. In our case, women had a median age 7 years older than men.

The prevalence of women with CS in the different published studies and registries ranges from 21% to 37%. (20-25) In our study it was 30%. Women were older and did not present more risk factors or comorbidities than men, as has been observed in some studies. (22-25)

Revascularization is the mainstay in the treatment of CS, (26) but some studies reveal considerable differences according to gender. (27) It even emerges from the analysis of the National (nationwide) Inpa-

Variable	Total (n=278)	Female (n=83)	Male (n=195)	p
Revascularization (CV)	222 (81)	65 (80)	157 (82)	0.768
Angina/ reAMI	9 (3)	4 (5)	5 (3)	0.331
Arrhythmias	88 (32)	26 (31)	62 (32)	0.939
AF	27(10)	6 (7)	21 (11)	0.362
VT/VF	39 (14)	11 (13,2)	28 (14,3)	0.808
AV block	16 (6)	7 (8)	9 (5)	0.211
Transient PM	15 (5)	2 (2)	13 (7)	0.151
ECV	36 (13)	7 (8)	29 (15)	0.143
Fever	36 (13)	11 (13)	25 (13)	0.922
Dialysis	20 (7)	6 (7)	14 (7)	0.988
Transfusion	21 (8)	7 (8)	14 (7)	0.717
Mortality	146 (53)	41(49)	105 (54)	0.470

AF: atrial fibrillation; CV: culprit vessel; ECV: electrical cardioversion; PM: pacemaker; reAMI: reinfarction; VT/VF: ventricular tachycardia/ventricular fibrillation  
Qualitative variables are presented as frequency and percentage.

**Table 2.** In-hospital evolution according to gender

	Dead (n= 146)	Alive (n= 132)	p
Age (years)	68 (60-78)	65 (57-76)	0.010
Female gender	41 (28)	42 (32)	0.470
HTN	113 (77)	93 (71)	0.223
Smoking	35 (24)	36 (28)	0.504
Type 2 diabetes	59 (40)	58 (44)	0.516
Previous AMI	30 (20)	14 (911)	0.025
STE-ACS	114 (78)	93 (71)	0.175
Revascularization	111 (77)	111 (87)	0.041
Swan Ganz	25 (17)	22 (17)	0.988
Balloon counterpulsation	37 (26)	24 (19)	0.170
MVD	106 (73)	91 (69)	0.551

AMI: acute myocardial infarction; HTN: Hypertension; MVD: Multiple vessel disease; STE-ACS: ST-segment elevation acute coronary syndrome  
Quantitative variables are presented as median and interquartile range, qualitative variables as frequency and percentage.

**Table 3.** Univariate analysis for mortality

tient Sample (NIS) database, with more than 134 000 elderly patients ( $\geq 75$  years) with CS, that women were less likely to receive angiography, angioplasty and mechanical support compared with men. (28) In our study there were no significant differences in revascularization between men and women, probably because most of the participating centers had available angioplasty and current guideline recommendations were followed, (29,30) but there was a difference in the indication of mechanical support, which was significantly lower in women, with the indication for balloon counterpulsation being half that of men. Extracorporeal membrane oxygenation was used in less than 4% of cases, which reflects the lack of availability and/or implementation given the socioeconomic conditions in Latin America. However, with the results of the IABP-SHOCK II trial, (31) those of ECMO-CS (32) and of ECLS-SHOCK, (33,34) which did not show benefits in survival, its routine use is not proposed for either men or women. The DanGer Shock study (35) in which Impella was used on a randomized basis, showed lower mortality with the use of the device in the included population, but in the subgroup analysis, only men benefited from its use. In Latin America, Impella was not used in any patient, although it should be noted that the registry was carried out prior to the publication of that study.

Mortality in our environment was high and there were no differences by gender as in other studies; (36,37) however, it is on this point where there is more controversy, since some recent studies report higher mortality in women. (38-40)

It should be noted that women may have some different baseline characteristics and that they are usually underrepresented in studies, but there have been no significant outcome differences with respect to gender, neither with revascularization (SHOCK trial), nor with the use of balloon counterpulsation (IABP-SHOCK II), or with revascularization of only the culprit vessel (CULPRIT-SHOCK). (41,42) There were also no differences according to gender in the use of other mechanical supports, except in the DanGer Shock study (35) in which the benefit of Impella was limited to the male group. In any case, it cannot be ruled out that this is due to the low number of women included. With the current data, both genders should continue to be treated equally. (43,44)

The heterogeneity of populations, treatment received and evolution makes it essential to have local and regional epidemiological data. Only by measuring and becoming aware of our own situation it is possible to evaluate whether it is necessary to implement change actions.

#### Limitations

The SCAI classification was not used, the registry did not report cardiac arrest at admission and there was no external audit of the data.

#### CONCLUSION

Latin American women who present with CS in the context of ACS are older than men, are revascularized in a similar manner, and have an equally high mortality. Although the use of mechanical supports was lower in women, this has not had an impact on mortality, nor has it had an impact on the results of randomized studies.

#### Acknowledgements

To the centers and researchers who participated in the registry: Hospital El Cruce, Argentina: Dr Mariano Adamowski; Hospital Nacional Edgardo Rebagliati Martins, Peru: Dr Maicol Cortez Sandoval; Hospital Luis Vernaza, Ecuador: Dr Freddy Pow Chon Long; Instituto Cabral de Corrientes, Argentina: Dr Stella Macin; Hospital de especialidades Dr Teodoro Maldonado Carbo, Ecuador: Dr Julio Burgos Acosta, Instituto nacional cardiovascular INCOR, Peru: Dr Manuel Chacon; Sanatorio Guemes, Argentina: Dr Joaquin Perea, Hospital Cemesa, Honduras: Dr Francisco Somoza; Hospital Univalle Norte, Bolivia: Dr Claudia Paz Soldan Patiño; Instituto Cardiovascular Migone, Paraguay: Dr Miguel Quintana; Clinica Pasteur, Argentina: Dr Andres Perelmuter; Instituto nacional de cardiologia Prof Dr Juan Cattoni, Paraguay: Dra Fabiola Rodriguez Caballero; Hospital Descentralizado Rawson, Argentina: Dra Maria De Los Angeles Navarta Navarro; Hospital Santojanni, Argentina: Dra Gabriela Velasco; Clinica Santa Isabel, Argentina: Dra Yanina Castillo Costa, Dr Flavio Delfino; Sanatorio Anchorena San Martin, Argentina: Dr Juan Souto; Hospital Español de Mendoza, Argentina: Dr Valentina Rodriguez; Sanatorio Pasteur de Catamarca, Argentina: Dr Maria Pia Marturano; Sanatorio Sagrado Corazón, Argentina: Dr Diego Costa; Hospital de especialidades Eugenio Espejo, Ecuador: Dr Diego Rengifo Escobar; Clinica Modelo Lanus, Argentina: Dr Diego Novielli; Hospital de Clinicas, Argentina: Dr Sandra Swieszkowski; Sanatorio Bernal, Argentina: Dr Juan Kinaszuk; Clinica Bazterrica, Argentina: Dr Victor Mauro; Instituto Cardiovascular de Buenos Aires, Argentina: Dr Leonardo Seoane; Sanatorio Modelo de Quilmes, Argentina: Dr Adrian Hrabar; Hospital Obrero 2, Bolivia: Dr Ruben Coca Pozo, Hospital El Carmen, Argentina: Dr Jose Di Milta; Sanatorio Americano, Argentina: Dr Mario Cifardoni; Sanatorio Adventista, Argentina: Dr Ramiro Ayala; Hospital Obrero 1, Bolivia: Dr Yhaquelina Quispe Villca; Hospital Barros Luco Trudeau, Chile: Dr Ricardo Ramirez; Clinica Privada Mayo, Argentina: Dr Eduardo Quinteros; Fundacion Favaloro, Argentina: Dr Fabricio Procopio; Hospital Italiano Regional del Sur, Argentina: Dr Veronica Heredia; Hospital Santa Isabel de Hungria, Argentina: Dr Maria Victoria Haedo; Sanatorio Allende, Argentina: Dr Nicolas Zaderenko; Sanatorio Anchorena, Argentina: Dr Alessio Raffaelli; Sanatorio Finochietto, Argentina: Dr Diego Crippa; Caja Nacional de Salud, Bolivia: Dr Richard Urquieta Cayoja; Hospital Nacional Guillermo Almenara, Peru: Dr Ciro Barrantes

#### Conflicts of interest

None declared.

(See authors' conflict of interests forms on the web).

## REFERENCES

1. Lucà F, Abrignani MG, Parrini I, Di Fusco SA, Giubilato S, Rao CM, et al. Update on management of cardiovascular diseases in women. *J Clin Med* 2022;11:1176-207. <https://doi.org/10.3390/jcm11051176>
2. Arora S, Stouffer GA, Kucharska-Newton AM, Qamar A, Vaduganathan M, Pandrey A, et al. Twenty year trends and sex differences in young adults hospitalized with acute myocardial infarction. *Circulation* 2019;139:1047-56. <https://doi.org/10.1161/CIRCULATIONAHA.118.037137>
3. Redfors B, Angerås O, Råmunddal T, Petursson P, Haraldsson I, Dworeck C, et al. Trends in gender differences in cardiac care and outcome after acute myocardial infarction in western Sweden: a report from the Swedish Web-system for Enhancement of Evidence-based Care in Heart Disease Evaluated According to Recommended Therapies (SWEDEHEART). *J Am Heart Assoc* 2015; 4:e001995. <https://doi.org/10.1161/JAHA.115.001995>
4. Koeth O, Zahn R, Heer T, Bauer T, Juenger C, Klein B, et al. Gender differences in patients with acute ST-elevation myocardial infarction complicated by cardiogenic shock. *Clin Res Cardiol* 2009;98:781-6. <https://doi.org/10.1007/s00392-009-0080-7>
5. Henry TD, Tomey MI, Tamis-Holland JE, Thiele H, Rao SV, Menon V, et al. Invasive Management of Acute Myocardial Infarction Complicated by Cardiogenic Shock: A Scientific Statement From the American Heart Association. *Circulation* 2021;143:815-29. <https://doi.org/10.1161/CIR.0000000000000959>
6. Castillo Costa Y, Delfino F, Mauro V, D'Imperio H, Barrero C, Charask A, et al. Clinical characteristics and evolution of patients with cardiogenic shock in Argentina in the context of an acute myocardial infarction with ST segment elevation. Data from the nationwide ARGENT-AMI-ST Registry. *Curr Probl Cardiol* 2023;48:e101468. <https://doi.org/10.1016/j.cpcardiol.2022.101468>
7. Nguyen HL, Saczynski JS, Gore JM, Goldberg RJ. Age and sex differences in duration of prehospital delay in patients with acute myocardial infarction: a systematic review. *Circ Cardiovasc Qual Outcomes* 2010;3:82-92. <https://doi.org/10.1161/CIRCOUTCOMES.109.884361>
8. Alabas OA, Gale CP, Hall M, Rutherford MJ, Szummer K, Lawesson SS, et al. Sex differences in treatments, relative survival, and excess mortality following acute myocardial infarction: national cohort study using the SWEDEHEART registry. *J Am Heart Assoc* 2017;6:e007123. <https://doi.org/10.1161/JAHA.117.007123>
9. Jneid H, Fonarow GC, Cannon CP, Hernandez AF, Palacios IF, Maree AO, et al. Sex differences in medical care and early death after acute myocardial infarction. *Circulation* 2008;118:2803-10. <https://doi.org/10.1161/CIRCULATIONAHA.108.789800>
10. Panchoy SB, Shantha GPS, Patel T, Cheskin LJ. Sex differences in short-term and long-term all-cause mortality among patients with ST-segment elevation myocardial infarction treated by primary percutaneous intervention: a meta-analysis. *JAMA Intern Med* 2014;174:1822-30. <https://doi.org/10.1001/jamainternmed.2014.4762>
11. Hunziker L, Radovanovic D, Jeger R, Pedrazzini G, Cuculi F, Urban P, et al. AMIS Plus Registry Investigators are listed in alphabetic order with the names of the local principal investigators. Twenty-Year Trends in the Incidence and Outcome of Cardiogenic Shock in AMIS Plus Registry. *Circ Cardiovasc Interv* 2019;12:e007293. <https://doi.org/10.1161/CIRCINTERVENTIONS.118.007293>
12. Arnegard ME, Whitten LA, Hunter C, Clayton J-A. Sex as a biological variable: a 5-year progress report and call to action. *J Womens Health* 2020;29:858-64. <https://doi.org/10.1089/jwh.2019.8247>
13. Shields CA, Wang X, Cornelius DC. Sex differences in cardiovascular response to sepsis. *Am J Physiol Cell Physiol* 2023;324:458-66. doi: 10.1152/ajpcell.00134.2022.
14. Xu J, Tong L, Yao J, Guo Z, Lui KY, Hu X, et al. Association of sex with clinical outcome in critically ill sepsis patients: a retrospective analysis of the large clinical database MIMIC-III. *Shock* 2019;52: 146-51. <https://doi.org/10.1097/SHK.0000000000001253>
15. Kouvari M, Souliotis K, Yannakoulia M, Panagiotakos DB. Cardiovascular Diseases in Women: Policies and Practices Around the Globe to Achieve Gender Equity in Cardiac Health. *Risk Manag Healthc Policy* 2020;13:2079-94. <https://doi.org/10.2147/RMHP.S264672>
16. Castillo Costa Y, Delfino F, Mauro V, D'Imperio H, Adamowski M, Cortez Sandoval MA, et al. Cardiogenic shock in the context of acute coronary syndromes in Latin America ("LATIN Shock"). *Curr Probl Cardiol* 2024;49:e102745. <https://doi.org/10.1016/j.cpcardiol.2024.102745>
17. Walli-Attaei M, Joseph P, Rosengren A, Chow CK, Rangarajan S, Lear SA, et al. Variations between women and men in risk factors, treatments, cardiovascular disease incidence, and death in 27 high-income, middle-income, and low-income countries (PURE): a prospective cohort study. *Lancet* 2020;396:97-109. [https://doi.org/10.1016/S0140-6736\(20\)30543-2](https://doi.org/10.1016/S0140-6736(20)30543-2)
18. Fraga CL, Macedo FVB, Rocha ARL, Ferreira Filho DSG, Nascimento BR. Gender Equity in Access to Reperfusion in Acute Myocardial Infarction: Still A Long Way to Go. *Arq Bras Cardiol* 2021;116:704-5. <https://doi.org/10.36660/abc.20210082>
19. Isorni MA, Aissaoui N, Angoulvant D, Bonello L, Lemesle G, Delmas C, et al. Temporal trends in clinical characteristics and management according to sex in patients with cardiogenic shock after acute myocardial infarction: The FAST-MI programme. *Arch Cardiovasc Dis* 2018;111:555-63. <https://doi.org/10.1016/j.acvd.2018.01.002>
20. Møller JE, Engstrøm T, Jensen LO, Eiskjær H, Mangner N, Polzin A, et al. Microaxial flow pump or standard care in infarct-related cardiogenic shock. *N Engl J Med* 2024;390:1382-93. <https://doi.org/10.1056/NEJMoa2312572>
21. Hochman JS, Sleeper LA, Webb JG, Samborn TA, White HD, Talley JD, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock. *N Engl J Med* 1999;34:625-34. <https://doi.org/10.1056/NEJM199908263410901>
22. Fengler K, Fuernau G, Desch S, Eitel I, Neumann F-J, Olbrich H-G, et al. Gender differences in patients with cardiogenic shock complicating myocardial infarction: a substudy of the IABP-SHOCK II-trial. *Clin Res Cardiol* 2015;104:71-8. <https://doi.org/10.1007/s00392-014-0767-2>
23. Arnold JH, Perl L, Assali A, Codner P, Greenberg G, Samara A, et al. The Impact of Sex on Cardiogenic Shock Outcomes Following ST Elevation Myocardial Infarction. *J Clin Med* 2023;12: 6259-70. <https://doi.org/10.3390/jcm12196259>
24. Vallabhajosyula S, Ya'Qoub L, Singh M, Bell MR, Gulati R, Cheungpasitporn W, et al. Sex Disparities in the Management and Outcomes of Cardiogenic Shock Complicating Acute Myocardial Infarction in the Young. *Circ Heart Fail* 2020;13:e007154. <https://doi.org/10.1161/CIRCHEARTFAILURE.120.007154>
25. Fengler K, Fuernau G, Desch S, Eitel I, Neumann F-J, Olbrich H-G, et al. Gender differences in patients with cardiogenic shock complicating myocardial infarction: a substudy of the IABP-SHOCK II-trial. *Clin Res Cardiol* 2015;104:71-8. <https://doi.org/10.1007/s00392-014-0767-2>
26. Naidu SS, Baran DA, Jentzer JC, Hollenberg SM, Diepen SV, Basir MB, et al. SCAI SHOCK Stage Classification Expert Consensus Update: A Review and Incorporation of Validation Studies: This statement was endorsed by the American College of Cardiology (ACC), American College of Emergency Physicians (ACEP), American Heart Association (AHA), European Society of Cardiology (ESC) Association for Acute Cardiovascular Care (ACVC), International Society for Heart and Lung Transplantation (ISHLT), Society of Critical Care Medicine (SCCM), and Society of Thoracic Surgeons (STS) in December 2021. *J Am Coll Cardiol* 2022;79:933-46.
27. Yan I, Schrage B, Weimann J, Dabboura S, Hilal R, Beer BN, et al. Sex differences in patients with cardiogenic shock. *ESC Heart Fail* 2021;8:1775-83. <https://doi.org/10.1002/ehf2.13303>
28. Vallabhajosyula S, Vallabhajosyula S, Dunlay SM, Hayes SN, Best PJM, Brenes-Salazar JA, et al. Sex and Gender Disparities in the Management and Outcomes of Acute Myocardial Infarction-Cardiogenic Shock in Older Adults. *Mayo Clin Proc* 2020;95:1916-27. <https://doi.org/10.1016/j.mayocp.2020.01.043>
29. Werdan K, Buerke M, Geppert A, Thiele H, Zwissler B, Ruß M; et al. Infarction-Related Cardiogenic Shock- Diagnosis, Monitoring and Therapy—A German-Austrian S3 Guideline. *Dtsch Arztebl Int* 2021;118:88-95. <https://doi.org/10.3238/arztebl.m2021.0012>
30. van Diepen S, Thiele H. An overview of international cardiogenic shock guidelines and application in clinical practice.

- Curr Opin Crit Care 2019;25:365-70. <https://doi.org/10.1097/MCC.0000000000000624>.
31. Thiele H, Zeymer U, Thelemann N, Neumann FJ, Hausleiter J, Abdel-Wahab M y cols. IABP-SHOCK II Trial (Intraaortic Balloon Pump in Cardiogenic Shock II) Investigators; IABP-SHOCK II Investigators. Intraaortic Balloon Pump in Cardiogenic Shock Complicating Acute Myocardial Infarction: Long-Term 6-Year Outcome of the Randomized IABP-SHOCK II Trial. *Circulation* 2019;139:395-403. <https://doi.org/10.1161/CIRCULATIONAHA.118.038201>.
32. Ostadal P, Rokyta R, Karasek J, Kruger A, Vondrakova D, Janotka M, et al. Extracorporeal Membrane Oxygenation in the Therapy of Cardiogenic Shock: Results of the ECMO-CS Randomized Clinical Trial. *Circulation* 2023;147:454-64. <https://doi.org/10.1161/CIRCULATIONAHA.122.062949>.
33. Thiele H, Zeymer U, Akin I, Behnes M, Rassaf T, Mahabadi AA, et al. Extracorporeal Life Support in Infarct-Related Cardiogenic Shock. *N Engl J Med* 2023;389:1286-97. <https://doi.org/10.1056/NEJMoa2307227>.
34. Zeymer U, Freund A, Hochadel M, Ostadal P, Belohlavek J, Rokyta R, et al. Veno arterial extracorporeal membrane oxygenation in patients with infarct-related cardiogenic shock: an individual patient data meta-analysis of randomised trials. *Lancet* 2023;402:1338-46. [https://doi.org/10.1016/S0140-6736\(23\)01607-0](https://doi.org/10.1016/S0140-6736(23)01607-0).
35. Møller JE, Engstrøm T, Jensen LO, Eiskjær H, Mangner N, Polzin A, y cols. DanGer Shock Investigators. Microaxial Flow Pump or Standard Care in Infarct-Related Cardiogenic Shock. *N Engl J Med* 2024;390:1382-93. <https://doi.org/10.1056/NEJMoa2312572>.
36. Wong SC, Sleeper LA, Monrad ES, Menegus MA, Palazzo A, Dzavik V, et al. Absence of gender differences in clinical outcomes in patients with cardiogenic shock complicating acute myocardial infarction. A report from the SHOCK Trial Registry. *J Am Coll Cardiol*. 2001; 38:1395-401. [https://doi.org/10.1016/s0735-1097\(01\)01581-9](https://doi.org/10.1016/s0735-1097(01)01581-9)
37. Fengler K, Fuernau G, Desch S, Eitel I, Neumann FJ, Olbrich HG, et al. Gender differences in patients with cardiogenic shock complicating myocardial infarction: a substudy of the IABP-SHOCK II-trial. *Clin Res Cardiol* 2015;104:71-8. <https://doi.org/10.1007/s00392-014-0767-2>
38. Ya'qoub L, Lemor A, Dabbagh M, O'Neill W, Khandelwal A, Martinez SC, et al. Racial, ethnic, and sex disparities in patients with STEMI and cardiogenic shock. *JACC Cardiovasc Interv* 2021;14:653-60. <https://doi.org/10.1016/j.jcin.2021.01.003>
39. Elgendy IY, Wegermann ZK, Li S, Mahtta D, Grau-Sepulveda M, Smilowitz NR, et al. Sex differences in management and outcomes of acute myocardial infarction patients presenting with cardiogenic shock. *JACC Cardiovasc Interv* 2022;15:642-52. <https://doi.org/10.1016/j.jcin.2021.12.033>
40. Fisher T, Hill N, Kalakoutas A, Lahlou A, Rathod K, Proudfoot A, et al. Sex differences in treatments and outcomes of patients with cardiogenic shock: a systematic review and epidemiological meta-analysis. *Crit Care* 2024;28:192-206. <https://doi.org/10.1186/s13054-024-04973-5>
41. Thiele H, Akin I, Sandri M, Fuernau G, de Waha S, Meyer-Saraei R, et al. CULPRIT-SHOCK Investigators. PCI Strategies in Patients with Acute Myocardial Infarction and Cardiogenic Shock. *N Engl J Med* 2017;377:2419-32. <https://doi.org/10.1056/NEJMoa1710261>.
42. Rubini Gimenez M, Zeymer U, Desch S, de Waha-Thiele S, Ouarrak T, Poess J y cols. Sex-Specific Management in Patients With Acute Myocardial Infarction and Cardiogenic Shock: A Substudy of the CULPRIT-SHOCK Trial. *Circ Cardiovasc Interv* 2020;13:e008537. <https://doi.org/10.1161/CIRCINTERVENTIONS.119.008537>.
43. Sambola A, Halvorsen S, Adlam D, Hassager C, Price S, Rosano G, et al. Management of cardiac emergencies in women: a clinical consensus statement of the Association for Acute Cardiovascular Care (ACVC), the European Association of Percutaneous Cardiovascular Interventions (EAPCI), the Heart Failure Association (HFA), and the European Heart Rhythm Association (EHRA) of the ESC, and the ESC Working Group on Cardiovascular Pharmacotherapy. *Eur Heart J Open* 2024;4:oeae011. <https://doi.org/doi:10.1093/ehjopen/oeae011>.
44. Mehta LS, Beckie TM, DeVon HA, Grines CL, Krumholz HM, Johnson MN, et al. Acute myocardial infarction in women: a scientific statement from the American Heart Association. *Circulation* 2016;133:916-47. <https://doi.org/10.1161/CIR.0000000000000351>