

Cardiogenic Shock: Evolution According to Gender in Latin America

Shock cardiogénico: evolución conforme al sexo en América Latina

HOLGER THIELE¹

Cardiogenic shock is still the most important cause of death in patients hospitalized with acute myocardial infarction. The randomized SHOCK (SHould we emergently revascularize Occluded Coronaries for cardiogenic shock) trial set the basis for an early invasive management of these patients, (1) with subsequent increase in early revascularization strategies across the globe. This strategy has dramatically improved outcome and reduced in-hospital mortality of cardiogenic shock patients from the former 70-80% to nowadays 40-50%. (2)

Currently, only few large-scale landmark randomized controlled trials have been performed in the cardiogenic shock setting including the above-mentioned SHOCK trial in 1999, (1) the IABP-SHOCK II trial, (3-5) the CULPRIT-SHOCK trial, (6,7) the ECLS-SHOCK trial, (8) and the recent DanGer-Shock trial. (9) Accordingly, only few measures rely on strong clinical evidence in the treatment of cardiogenic shock. (10,11)

When insufficient evidence is available and mortality still high, evidence from observational data is important and additive to randomized data. This is particularly true for sex specific differences in cardiogenic shock because often women are underrepresented in the large-scale randomized controlled trials. This is partly an effect because women are older and many large-scale randomized controlled trials had an upper age limit for inclusion which automatically leads to less women for inclusion because of the higher age in women presenting with cardiogenic shock. Therefore, it is even more important in clinical practice to measure the outcome of acute coronary syndromes and the complications including cardiogenic shock with respect to sex specific differences. Only by measuring outcome, measures can be implemented to improve outcome for women and also men. As such, it can only be supported to see the publication of the

LATIN Shock registry from Argentina, Bolivia, Chile, Ecuador, Honduras, Paraguay and Peru. (12) The in-hospital mortality of 49% in women and 54% in men shows the still very high mortality in cardiogenic shock in the current era of early revascularization. (13) Interestingly, despite the older age in women there was no difference in mortality, which has also been shown in other analyses such as the CULPRIT-SHOCK and IABP-SHOCK II sex specific subanalyses. (14,15) In contrast, other observational data suggest less invasive treatment in females presenting with cardiogenic shock with subsequent higher mortality. (16) Accordingly, the LATIN Shock registry supports no relevant outcome differences in cardiogenic shock based on sex.

Interestingly, still the majority of patients is treated by intraaortic balloon pumping where the evidence does not support to use this device. (3-5) On the other hand evidence for active mechanical circulatory support is also limited and currently only the DanGer-Shock trial and a meta-analysis of all trials comparing active mechanical circulatory support versus control supports the use in very selected patients with ST-elevation myocardial infarction and no risk of hypoxic brain injury. (9,17)

The authors should be congratulated to put this LATIN Shock registry together. More efforts should be directed towards cardiogenic shock registries and a higher number of patients will help to define the best treatment strategies to improve outcome in cardiogenic shock also with respect to sex specific differences.

Ethical considerations

Not applicable.

Conflicts of interest

None declared.

(See authors' conflict of interests forms on the web).

REV ARGENT CARDIOL 2024;92:399-400. <http://dx.doi.org/10.7775/rac.v92.i6.20846>

SEE RELATED ARTICLE: Rev Argent Cardiol 2024;92:403-409. <http://dx.doi.org/10.7775/rac.v92.i6.20842>

Correspondence: Holger Thiele, MD. Director. Heart Center Leipzig at Leipzig University. Department of Internal Medicine/Cardiology Strümpellstr. 39. D-04289 Leipzig. Germany. e-mail: holger.thiele@medizin.uni-leipzig.de



<https://creativecommons.org/licenses/by-nc-sa/4.0/>

©Revista Argentina de Cardiología

¹ Heart Center Leipzig at Leipzig University, Leipzig, Germany

REFERENCES

1. Hochman JS, Sleeper LA, Webb JG, Sanborn TA, White HD, Talley JD, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK Investigators. Should We Emergently Revascularize Occluded Coronaries for Cardiogenic Shock. *N Engl J Med* 1999;341:625-34. <https://doi.org/10.1056/NEJM199908263410901>
2. Thiele H, Ohman EM, Desch S, Eitel I, de Waha S. Management of cardiogenic shock. *Eur Heart J* 2015;36:1223-30. <https://doi.org/10.1093/eurheartj/ehv051>
3. Thiele H, Zeymer U, Neumann F-J, Ferenc M, Olbrich H-G, Hausleiter J, et al. Intraaortic balloon counterpulsation in acute myocardial infarction complicated by cardiogenic shock. Final 12-month results of the randomised IntraAortic Balloon Pump in cardiogenic shock II (IABP-SHOCK II) Trial. *Lancet*. 2013;382:1638-45. [https://doi.org/10.1016/S0140-6736\(13\)61783-3](https://doi.org/10.1016/S0140-6736(13)61783-3)
4. Thiele H, Zeymer U, Neumann F-J, Ferenc M, Olbrich H-G, Hausleiter J, et al. Intraaortic balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med*. 2012;367:1287-96. <https://doi.org/10.1056/NEJMoa1208410>
5. Thiele H, Zeymer U, Thelemann N, Neumann F-J, Hausleiter J, Abdel-Wahab M, et al. Intraaortic balloon pump in cardiogenic shock complicating acute myocardial infarction. Long-term 6-year outcome of the randomized IABP-SHOCK II Trial. *Circulation*. 2019;139:395-403. doi: <https://doi.org/10.1161/CIRCULATIONAHA.118.038201>
6. Thiele H, Akin I, Sandri M, de Waha-Thiele S, Meyer-Saraei R, Fuernau G, et al. One-year outcomes after PCI strategies in cardiogenic shock. *N Engl J Med* 2018;379:1699-710. <https://doi.org/10.1056/NEJMoa1808788>
7. Thiele H, Akin I, Sandri M, Fuernau G, de Waha S, Meyer-Saraei R, et al. PCI strategies in patients with acute myocardial infarction and cardiogenic shock. *N Engl J Med* 2017;377:2419-32. <https://doi.org/10.1056/NEJMoa1710261>
8. Thiele H, Zeymer U, Akin I, Behnes M, Rassaf T, Mahabadi A, et al. Extracorporeal life support in infarct-related cardiogenic shock. *N Engl J Med* 2023;389:1286-97. <https://doi.org/10.1056/NEJMoa2307227>
9. Møller JE, Engstrøm T, Jensen LO, Eiskjær H, Mangner N, Polzin A, Schulze PC, Skurk C, Nordbeck P, Clemmensen P, et al. Microaxial flow pump or standard care in infarct-related cardiogenic shock. *N Engl J Med* 2024;390:1382-93. <https://doi.org/10.1056/NEJMoa2312572>
10. Thiele H, Ohman EM, de Waha-Thiele S, Zeymer U, Desch S. Management of cardiogenic shock complicating myocardial infarction: an update 2019. *Eur Heart J* 2019;40:2671-83. <https://doi.org/10.1093/eurheartj/ehz363>
11. Zeymer U, Bueno H, Granger CB, Hochman J, Huber K, Lettino M, et al. ACCA - Position paper for the diagnosis and treatment of patients with acute myocardial infarction complicated by cardiogenic shock. *Eur Heart J Acute Cardiovasc Care* 2020;9:183-97. <https://doi.org/10.1177/2048872619894254>
12. Castillo Costa Y, Delfino F, Mauro V, D Imperio H, Adamowski M, Cortez Sandoval MA, et al. Cardiogenic shock in the context of acute coronary syndromes in Latin America ("LATIN Shock"). *Curr Probl Cardiol*. 2024;49:102745. <https://doi.org/10.1016/j.cpcardiol.2024.102745>
13. Castillo Costa Y, Delfino F, Macías J, Quintana M, Adamowski M, Rodríguez Caballero F, et al. Characteristics and Evolution of Cardiogenic Shock According to gender in Latin America. *LATIN Shock Registry Data*. *Rev Argent Cardiol* 2024;92:403-9. <http://dx.doi.org/10.7775/rac.v92.i6.20842>
14. Rubini Gimenez M, Zeymer U, Desch S, de Waha-Thiele S, Ouararak T, Poess J, Meyer-Saraei R, Schneider S, Fuernau G, Stepinska J, et al. Sex-specific management in patients with acute myocardial infarction and cardiogenic shock. *Circulation: Cardiovasc Interv* 2020;13:e008537. doi: <https://doi.org/10.1161/CIRCINTERVENTIONS.119.008537>
15. Fengler K, Fuernau G, Desch S, Eitel I, Neumann FJ, Olbrich HG, de Waha A, de Waha S, Richardt G, Hennersdorf M, et al. Gender differences in patients with cardiogenic shock complicating myocardial infarction: a substudy of the IABP-SHOCK II-trial. *Clin Res Cardiol* 2015;104:71-8. <https://doi.org/10.1007/s00392-014-0767-2>
16. Vallabhajosyula S, Ya'Qoub L, Singh M, Bell MR, Gulati R, Cheungpasitporn W, et al. Sex disparities in the management and outcomes of cardiogenic shock complicating acute myocardial infarction in the young. *Circ Heart Fail* 2020;13:e007154. <https://doi.org/10.1161/circheartfailure.120.007154>
17. Thiele H, Møller JE, Henriques JPS, Bogerd M, Seyfarth M, Burkhoff D, Ostadal P, Rokyta R, Belohlavek J, Massberg S, et al. Temporary mechanical circulatory support in infarct-related cardiogenic shock. An individual patient data meta-analysis of all randomised trials with 6-month follow-up. *Lancet* 2024;404:1019-28. [https://doi.org/10.1016/S0140-6736\(24\)01448-X](https://doi.org/10.1016/S0140-6736(24)01448-X)