

Cocaine and Cocaine Base Paste in Cardiology. A matter to Consider

Cocaína y paco en cardiología. Un tema que considerar

ROCÍO CASTILLA¹, FRANCISCO MATHEU², FRANCISCO AZZATO², JOSÉ MILEI¹

INTRODUCTION

Twenty years ago, we published in this journal a typical case of cocaine-induced dilated cardiomyopathy. Cocaine (benzoylecgonine) is a natural alkaloid extracted from the leaves of the *Erythroxylum coca* bush. (1) This condition is frequently observed in persons addicted to cocaine, and in that case report we demonstrated the presence of baseline hypokinesia that responded to dobutamine stress echocardiography, with angiographically normal coronary arteries and pathognomonic anatomic-pathological study. We found loss of the sarcomeric structure coupled with deep mitochondrial abnormalities and increased interstitial fibrosis without lymphocyte infiltration. (1) Coronary artery spasm and sudden cardiac death are common complications of cocaine abuse. By that time, we called the attention of the press to the effects of the “damned cocaine”. (5)

Drug abuse is an extremely serious problem worldwide. The 2013 World Drug Report of the United Nations Office on Drugs and Crime (UNODC) places Argentina among the countries where cocaine use among the population between 15 and 64 years is between 0.51% and 1%. Furthermore, trends in cocaine users by annual prevalence in Latin America and the Caribbean increased from 15% to 21% between 2004-2005 and 2011. (6)

A recent report by the Drug Abuse Prevention and Fight against Drug Trafficking Programming Secretariat (Secretaría de Programación para la Prevención de Drogadicción y Lucha contra el Narcotráfico, SEDRONAR) in 2017 through the Argentine Drug Observatory National Directory (Dirección Nacional del Observatorio Argentino de Drogas, DNOAD) indicates that 5.3% of the population between 12 and 65 years old has used cocaine at some time in their lives, representing more than one million people. (7) In addition, out of the 292,052 persons who used cocaine during the period 2016-2017, 105,028 presented drug dependence. An alarming data is that among them, 2,165 were children and adolescents. (8)

In turn, in 2007, the SEDRONAR had already reported an increase of more than 200% in the use of cocaine base paste or paco in our country. (9)

Effectively, cocaine base paste, known by its acronym paco (“Pasta base de COcaína”), is a relatively new drug. Cocaine base paste is the intermediate product in the process of extraction and purification of cocaine hydrochloride, in which organic solvents as toluene or kerosene and other products, as sulfuric acid, potassium permanganate or hydrochloric acid are used. (10)

Cocaine base paste is a “smokable cocaine”. It is smoked alone or mixed with tobacco, marijuana or both.

The product obtained is then adulterated by “retailers”. (11) Inactive adulterants as talc or lactose are usually added as fillers and active adulterants can also be used to increase or mimic the effect of the drug. Stimulants as caffeine or amphetamines are mixed with cocaine to increase addiction, or with compounds with anesthetic effect, as lidocaine, etc.

The quality of cocaine base paste varies according to the substances used to obtain the drug and to the adulterants, dissimilar substances with diverse toxic effects, all included under the term “paco”.

Analytical data

In our country, the chair of Toxicology and Legal Chemistry, Analytical Toxicology Advise Laboratory (CENATOXA), (12-16) has extensive experience in the determination of abuse drugs, especially cocaine. The laboratory has developed analytical methods for the investigation of these drugs, both by immunological methods or gas chromatography combined or not with mass spectrometry (GC-MS). Twenty-five samples of cocaine commercialized as cocaine base paste were analyzed in this laboratory. A qualitative analysis was carried out following a similar protocol to that recommended by the United Nations.

The content of cocaine showed great variability even though the average purity was > 50% in most of

REV ARGENT CARDIOL 2020;88:452-454. <http://dx.doi.org/10.7775/rac.v88.i5.18805>

Address for reprints: José Milei - IATIMET UBA-CONICET - M.T. de Alvear 2270 - 1115-CABA - E-mail: josemillei@gmail.com

¹ CONICET, University of Buenos Aires, Instituto Alberto C. Taquini de Investigaciones en Medicina Traslacional (IATIMET), School of Medicine, Buenos Aires, Argentina.

² University of Buenos Aires, School of Medicine, Sixth Chair of Medicine. Hospital de Clínicas. Buenos Aires, Argentina

the samples. Cocaine was the basic drug found in all the samples analyzed with relative concentrations between 28.7% and 89.0% of purity (mean purity 66.24% and median purity 64.6%).

The analytical data revealed the presence of different alkaloids, either natural or those obtained during the process of cocaine production or by cocaine degradation or hydrolysis. The alkaloids found, expressed as a percentage of samples, were cocaine (100%), cis/trans-cinnamoylcocaine (100%), tropacocaine (100%), anhydromethylecgonine (100%), methylecgonine (100%), norcocaine (100%) and benzoylecgonine (32%).

Caffeine was the only adulterant found, with relative concentrations between 21.7% and 23.4%. In another study carried out by SEDRONAR in 2016 to analyze the composition of cocaine base paste in samples seized in Argentina, lidocaine and phenacetin were found in 32% and 29% of the samples, respectively, in addition to caffeine. (17) The presence of these adulterants demonstrates that cocaine base paste users are exposed to more than one stimulant and addictive substance. (18)

At the same time, the CENOTOXA laboratory detected the presence of Diuron [3-(3,4-dichlorophenyl)-1,1-dimethylurea] in 28% of the samples, a urea-derived herbicide used to spray the coca leaves. The presence of this herbicide demonstrates that other substances capable of affecting health can be present in cocaine base paste.

It is necessary to continue research to accurately determine the composition of the product commercialized as paco to contribute to a better evaluation of the nature of the emerging problem in health, social and control areas.

Effects of cocaine base paste

Cocaine base paste contains highly liposoluble alkaloids that quickly pass the blood-brain barrier and reach the central nervous system in five seconds. This is one of the reasons for its great addictive power.

The effects of cocaine base paste are categorized in the following stages: (11)

1. Euphoria: sense of wellbeing, lessening of inhibition, sense of pleasure, ecstasy, changes in attention levels, hyperexcitability, sense of competence and capability, accelerated thinking processes, lessening of hunger and fatigue, hypervigilance, and increased blood pressure, body temperature and respiratory rate.
2. Dysphoria: anxiety, depression and instability, uncontrollable desire to continue smoking, sadness, apathy and sexual indifference.
3. Uninterrupted use, even with blood levels enough to prevent the stage of dysphoria.
4. Psychosis and hallucinations: psychosis or loss of contact with reality, which can occur after several days or weeks of frequent smoking. This stage can last weeks or months. Hallucinations can be visual, auditory, olfactory or tactile.

Other effects of chronic use of cocaine base paste are hypertension, tachycardia, weight loss, pallor, insomnia, verbiage, mydriasis, nausea or vomiting, dry mouth, tremor, lack of coordination, headache, dizziness and itching. (11)

When use is stopped, a severe withdrawal syndrome develops, characterized by severe depression, irritability and suicidal ideation.

In spite of being an intermediate product of cocaine and that many of its effects are related to cocaine content, the clinical manifestations of cocaine base paste are different from those observed with cocaine hydrochloride, possibly due to the impurity of this drug. (11) However, considering that users usually use two drugs or more, cardiovascular involvement cannot be ruled out. In this sense, an adequate cardiological examination cannot be absent.

Main contributions of the bibliography

There are few experimental scientific publications in the international literature about cocaine base paste (19) and they mention that the different solvents and adulterants determine different "qualities" of the drug called paco, which can contain very different products with different toxic effects.

The most encountered respiratory symptoms include cough productive of black pigmented-sputum, dyspnea and bronchospasm. Chest X-rays have shown bilateral radiopaque perihilar infiltrates and, in some cases, nodular airspace opacities with areas of air trapping. (21)

Pulmonary lesions due to cocaine base paste are well documented, and include pneumomediastinum, (22), pulmonary edema (23-26), pulmonary dysfunction (27) and decreased diffusing capacity. (28)

Electrocardiographic abnormalities are common with cocaine base paste and have been reported in 60% of users, with 41% long QTc cases. Other abnormalities include incomplete right bundle branch block, junctional rhythm, and signs of ischemia or necrosis that were more common than in the control group. (29)

Previgliano et al. (30) demonstrated global cerebral hypoperfusion associated with increased vascular resistance in the anterior cerebral circulation in chronic users, which could be attributed to the sympathomimetic properties of cocaine. (31)

Furthermore, although there are post-mortem specimens of drug abusers, the findings are contaminated by the coexistence of multiple toxic agents (paco, alcohol, cocaine, marijuana, tobacco, sedatives, hypnotics, etc.) in the same patient. That is why it is extremely important to know the effects on health of cocaine base paste in the absence of other toxic substances. For this reason, studies have been conducted on animal models. As we have previously mentioned, there are not many scientific publications on the matter, and in such investigations the drug was administered using different routes than those used by drug

users; thus, the results obtained cannot be extrapolated to humans.

CONCLUSION

In spite of being an intermediate product of cocaine and that many of its effects are related to cocaine content, the clinical manifestations of cocaine base paste are different from those observed with cocaine hydrochloride, probably due to the impurity of this drug. Although there are post-mortem specimens of drug abusers, the findings are contaminated by the coexistence of multiple toxic agents in the same patient. Despite the great social impact produced by the use of this drug, little is known about its chemical composition, chronic effects on the different organs or pathophysiology. The creation of multidisciplinary groups, with modern technology and adequate economic support is urgent to deal with this extremely serious scourge. In that sense, the cardiologist approach in these patients cannot be absent.

Conflicts of interest

None declared.

(See authors' conflicts of interest forms on the website/ Supplementary material)

Acknowledgements

This work was supported by PIO CONICET- SEDRONAR 2015-2016- 15120150100007CO

Ethical considerations:

Not applicable

REFERENCES

- Álvarez CB, Pettinari M, Nijensohn S, Killinger C, Milei J, Chiale P. Dilated Myocardiopathy due to cocaine abuse. *Rev Argent Cardiol* 2000;68:729-34.
- Manninger M, Perl S, Brussee H, Toth GG. Sniff of coke breaks the heart: cocaine-induced coronary vasospasm aggravated by therapeutic hypothermia and vasopressors after aborted sudden cardiac death: a case report. *Eur Heart J - Case Reports* 2018;2:1-4. <https://doi.org/10.1093/ehjcr/ty041>
- Priori SG, Blomstrom-Lundqvist C. European Society of Cardiology Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death summarized by co-chairs. *Eur Heart J* 2015;36:2757-9. <https://doi.org/10.1093/eurheartj/ehv445>
- Lucena J, Blanco M, Jurado C, Rico A, Salguero M, Vázquez R, et al. Cocaine-related sudden death: a prospective investigation in south-west Spain. *Eur Heart J* 2010;31:318-29. <https://doi.org/10.1093/eurheartj/ehp557>
- Milei J. La cocaína castiga al corazón. Sección Opinión. Edición del 3 de febrero de 2000. Clarín. Buenos Aires.
- United Nations Office on Drugs and Crime (UNODC) World Drug Report 2013. Vienna. United Nations, New York.
- DNOAD. Estudio nacional en población de 12 a 65 años, sobre consumo de sustancias psicoactivas. Argentina, 2017. Informe de resultados N.º 1: Magnitud del consumo de sustancias a nivel nacional.
- Secretaría de Programación para la Prevención de Drogadicción y Lucha contra el Narcotráfico Resumen de los resultados del estudio 2017 de consumo de sustancias psicoactivas.
- Informe Junta Internacional de Fiscalización de Estupefacientes (JIFE) (2007) – Organización de las Naciones Unidas (ONU).
- Clandestine manufacture of substances under international control. Manual for use by national law enforcement authorities and personnel of narcotics laboratories. Division of narcotic drugs Vienna. United Nations. New York, 1987.
- Castilla R, Matheu F, Azzato F, Milei J. Productos intermedios de la cocaína: la pasta base o "paco". *AMA* 2020. [En prensa].
- Quiroga PN, Olivera NM, Vignati K, Ridolfi A, Villamil Lepori EC. Estudio de los compuestos presentes en muestras no biológicas de cocaína analizadas en el CENATOXA. *Acta Toxicol Argent* 2009;17:24.
- Olivera NM, Jerez GA, Vignati KG, Ridolfi A, Villamil Lepori EC, Quiroga P, et al. Contenido de alcaloides en muestras de paco analizadas en CENATOXA. *Acta Toxicol Argent* 2010;18:32.
- Jerez GA, Olivera NM, Vignati KG, Ridolfi A, Villamil Lepori EC, Quiroga P. Concentración de cocaína y cafeína en muestras comercializadas bajo la denominación de "paco" analizadas en CENATOXA. *Acta Toxicol Argent* 2010;18:28.
- Olivera NM, Jerez GA, Vignati KG, Ridolfi A, Villamil Lepori EC, Quiroga P. Perfil de alcaloides y concentración de cocaína en muestras de cocaína base analizadas en CENATOXA. *Ciencia Forense Latinoamericana* 2011;3:23.
- Quiroga P, Olivera NM, Jerez GA, Vignati KG. Estudio analítico de muestras de paco. *Acta Toxicol Argent* 2012;20:22.
- La Comisión Interamericana para el Control del Abuso de Drogas (CICAD), Observatorio Inter-Americano sobre Drogas (OID). Análisis de la caracterización química de cocaínas fumables 2016.
- Prieto JP, Galvalisi M, López-Hill X, Meikle MN, Abin-Carriquiry JA, Scorza C. Caffeine enhances and accelerates the expression of sensitization induced by coca paste indicating its relevance as a main adulterant. *Am J Addict* 2015;24:475-81. <https://doi.org/10.1111/ajad.12245>
- Umpierrez E (2006). Informe del laboratorio de análisis orgánico, Departamento de Química Orgánica, Facultad de Química – Polo tecnológico. Informe 06-17 JND- INFO.
- Pascale A, Negrín A, Ormaechea R. Preliminary study of the effect on the lungs due to the consumption of cocaine base. *Arch Bronconeumol* 2011;47:108. <https://doi.org/10.1016/j.arbres.2010.08.007>
- Kline JN, Hirasuna JD. Pulmonary edema after freebase cocaine smoking-not due to an adulterant. *Chest* 1990;97:1009-10. <https://doi.org/10.1378/chest.97.4.1009>
- Sullivan TP, Pierson DJ. Pneumomediastinum after freebase cocaine use. *AJR Am J Roentgenol* 1997;168:84. <https://doi.org/10.2214/ajr.168.1.8976925>
- Laposata EA, Mayo GL. A review of pulmonary pathology and mechanisms associated with inhalation of freebase cocaine ("crack"). *Am J Forensic Med Pathol* 1993;14:1-9. <https://doi.org/10.1097/0000433-199303000-00001>
- Cucco RA, Yoo OH, Cregler L, Chang JC. Nonfatal pulmonary edema after "freebase" cocaine smoking. *Am Rev Respir Dis* 1987;136:179-81. <https://doi.org/10.1164/ajrccm/136.1.179>
- Allred RJ, Ewer S. Therapy for pulmonary edema following IV "freebase" cocaine use. *Ann Emerg Med* 1982;11:228-30. [https://doi.org/10.1016/S0196-0644\(82\)80507-6](https://doi.org/10.1016/S0196-0644(82)80507-6)
- Allred RJ, Ewer S. Fatal pulmonary edema following intravenous "freebase" cocaine use. *Ann Emerg Med* 1981;10:441-2. [https://doi.org/10.1016/S0196-0644\(81\)80314-9](https://doi.org/10.1016/S0196-0644(81)80314-9)
- Itkonen J, Schnoll S, Glassroth J. Pulmonary dysfunction in 'freebase' cocaine users. *Arch Intern Med* 1984;144:2195-7. <https://doi.org/10.1001/archinte.1984.04400020107016>
- Weiss RD, Tilles DS, Goldenheim PD, Mirin SM. Decreased single breath carbon monoxide diffusing capacity in cocaine freebase smokers. *Drug Alcohol Depend* 1987;19:271-6. [https://doi.org/10.1016/0376-8716\(87\)90047-0](https://doi.org/10.1016/0376-8716(87)90047-0)
- Aguilera X, Arribada A, Apt W, Rodríguez J, Aqueveque A. Electrocardiographic abnormalities in cocaine base paste consumers of the metropolitan area. *Rev Med Chil* 1997;125:143-7.
- Previgliano I, Cortese S, Di Nardo V, et al. Changes in cerebral hemodynamics in chronic users of "PACO" and cocaine: case-control study. *Vertex* 2014;25:92-8.
- López-Hill A, Prieto JP, Meikle MN, Urbanavicius J, Abin-Carriquiry JA, Prunell G, et al. Coca-paste seized samples characterization: chemical analysis, stimulating effect in rats and relevance of caffeine as a major adulterant. *Behav Brain Res* 2011;221:134-41. <https://doi.org/10.1016/j.bbr.2011.03.005>